

**THE SPERMATOGENESIS  
OF ANASA TRISTIS, PP.  
223 - 272**

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649310258

The Spermatogenesis of Anasa Tristis, pp. 223 - 272 by F. C. Paulmier

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd.  
Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

[www.triestepublishing.com](http://www.triestepublishing.com)

**F. C. PAULMIER**

**THE SPERMATOGENESIS  
OF ANASA TRISTIS, PP.  
223 - 272**



THE  
SPERMATOGENESIS OF ANASA TRISTIS

SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS  
FOR THE DEGREE OF DOCTOR OF PHILOSOPHY, IN THE  
FACULTY OF PURE SCIENCE, COLUMBIA UNIVERSITY

*Fredrick  
C. Paulmier*  
F. C. PAULMIER

---

Reprinted from JOURNAL OF MORPHOLOGY, Supplement to Vol. XV.

---

BOSTON, U.S.A.  
GINN & COMPANY, PUBLISHERS  
*The Athenæum Press*  
1899

FREDERICK CLARK PAULMIER.

*Born Madison, New Jersey, May 17, 1873.*

GRADUATE MORRIS ACADEMY, MORRISTOWN, N.J., JUNE, 1890.

B.S., PRINCETON UNIVERSITY, JUNE, 1894.

M.S., PRINCETON UNIVERSITY, JUNE, 1896.

CLASS OF '77 UNIVERSITY FELLOW IN BIOLOGY, PRINCETON UNIVERSITY, 1894-95.

GRADUATE STUDENT, DEPT. OF ZOOLOGY, COLUMBIA UNIVERSITY, 1895-98.

FELLOW IN ZOOLOGY, COLUMBIA UNIVERSITY, 1898-99.

ASSISTANT IN ZOOLOGY, COLUMBIA UNIVERSITY, 1899-

---

PREVIOUS PUBLICATION.

Chromatin Reduction in the Hemiptera.

*Anat. Anz.*, Bd. XIV, 1898.



# THE SPERMATOGENESIS OF ANASA TRISTIS.

F. C. PAULMIER.

CONTENTS.		PAGE
INTRODUCTION .....		224
I. MATERIAL AND METHODS.....		226
II. DESCRIPTION OF TESTIS.....		226
III. THE SPERMATOGONES.....		228
Food Cells.....		231
IV. THE MATURATION DIVISIONS.....		232
1. Origin of the Tetrads .....		232
<i>a.</i> Early History of the Chromatin.....		232
<i>b.</i> Formation of the Tetrads.....		233
<i>c.</i> The Small Chromosomes.....		235
<i>d.</i> The Achromatic Structures.....		236
2. The First Division.....		237
<i>a.</i> The Tetrads .....		237
<i>b.</i> The Achromatic Structures.....		239
<i>c.</i> Separation of the Chromatin Elements.....		241
3. The Second Division.....		243
<i>a.</i> Achromatic Structures.....		243
<i>b.</i> The Chromatin.....		243
<i>c.</i> Later History of the Achromatic Structures.....		244
V. FORMATION OF THE SPERMATOZOÏN.....		247
<i>a.</i> The Nucleus.....		247
<i>b.</i> The Nebenkern.....		247
<i>c.</i> The Centrosome.....		249
<i>d.</i> The Acrosome .....		250
<i>e.</i> Further Development of the Spermatozoa.....		251
<i>f.</i> Giant Spermatozoa.....		254
VI. REVIEW OF TETRAD FORMATION AND SMALL CHROMOSOMES.....		256
<i>a.</i> Tetrad Formation.....		256
<i>b.</i> The Small Chromosomes.....		259
VII. GENERAL SUMMARY.....		263



## INTRODUCTION.

THE following paper contains the results of a study of the male germ cells of *Anasa*, one of the Heteroptera, corroborating in the main an earlier work by Henking on *Pyrrhocris*, although a number of new points are brought out to supplement the excellent work of my predecessor.<sup>1</sup>

Henking ('91) described the structure of the spermatogones and the preparation of the chromatin for the spermatocyte divisions, and though many of the details escaped him, he certainly, it seems, made a correct estimate of the value of these divisions. His description of the formation of the spermatid is especially good, and with the exception of what are probably merely specific variations, the process in *Anasa* approximates very closely to what he has described. Henking was also the first to describe the peculiar chromatic body in the spermatocytes to which particular attention has been paid in this paper under the name of the small chromosome.

Wilcox ('95, '96), the only author besides Henking who has followed the entire history of the spermatozoa in the insects, found that in *Caloptenus*, one of the Orthoptera, the process varies in many essentials from that in the Heteroptera, the most fundamental difference lying in the fact that he finds no longitudinal splitting in the spermatocytes; and although tetrads are formed (by conjugation), both subsequent divisions are transverse or "reducing" ones.

Besides these, several authors have given a partial account of the history of the spermatozoa. Perhaps the most important of these works is Vom Rath's ('92) on *Gryllotalpa*, where the chromatin changes, and especially the process of reduction,

<sup>1</sup> A considerable number of observations have been made on other species of Hemiptera-Heteroptera, the results of some of which have been already described (Paulmier, '98). Among the forms examined have been species of *Euchistus*, *Brochymena*, *Mormidea*, *Murgantia*, *Coenus*, *Podisus*, *Mineus*, and *Neara* of the family Pentatomidae; *Chariesterus*, *Euthocta*, and *Alydus* of the family Coreidae, and *Myodocha* and *Lygaeus* of the family Lygaeidae. From these observations it appears that there may be a very great difference in details between the various genera (cf. difference in tetrad formation described before, '98), but the main results are the same in all cases. Some observations on the spermatogenesis of *Papilio* gave interesting results which will be mentioned in the text.

were first carefully worked out. Platner ('89), on the other hand, confined himself to a study of the achromatic structures in the Lepidoptera.

Montgomery in three papers ('97-'99) has treated of the spermatogenesis of *Pentatoma (Euchistus)*, one of the Heteroptera, carrying it, however, only up to the formation of the spermatid. His results differ in many respects from mine; in regard to the reduction question his conclusion is that while a longitudinal splitting of the chromosomes and consequently an equatorial division usually occurs in the spermatocyte divisions, both divisions may in some cases be transverse, and therefore both reducing. He gives an interesting account of the body which he calls the "chromatin nucleolus," and which I have described under the name of the small chromosome.

In the present work I have endeavored to follow the entire history of the male germ cells, from the spermatogones to the mature spermatozoa. Among the more important results may be mentioned the probable continuity of the centrosome throughout all stages, of interest on account of its bearing on the discussion as to whether or not the centrosome is a permanent organ of the cell; the occurrence of a longitudinal splitting of the chromosomes in the prophase of the first spermatocyte division and its relation to the formation of tetrads, a question of considerable interest in view of several works on insect spermatogenesis, and in which my observations point to an agreement with the copepod type of tetrad formation; the history of a chromatic body which has been described recently by several authors, and a suggestion as to its meaning, which is of interest as bearing on the theory of the individuality of the chromosomes; and a partial explanation of the giant spermatozoa.

I wish to express my best thanks to Professor Wilson for his advice and encouragement throughout the progress of this work. My thanks are also due to Dr. Calkins for his kindness in reading the manuscript of this paper.

## I. MATERIAL AND METHODS.

The insect on which most of the following observations were made is *Anasa tristis*, De Geer, of the family Coreina, order Hemiptera-Heteroptera. It is commonly known as the "squash bug," its favorite food plant being the squash, though it also feeds on the pumpkin and probably other plants of the same family. The adult bugs winter under the bark of trees and under stones or in crevices, and appear during the latter part of June and July, the eggs being laid during the latter month. The larval period lasts about four weeks, and abundant material for spermatogenesis can be obtained in any desired stage. Most of the material used was obtained from specimens of the two instars preceding the last moult, the majority of the cells of the testis then being in the spermatocyte division or early spermatid stage. The later growth of the spermatozoa takes place after the last moult and during the winter.

The testes were dissected out in normal salt solution and brought as rapidly as possible into the fixing fluid. For fixing, a number of agents were tried, but the best results were obtained with Hermann's and Fleming's fluids, used for fifteen to thirty minutes and followed by thorough washing in running water. The material was then gradually transferred through successive alcohols to 90 per cent, in which it was kept. It was cleared in clove oil and imbedded in paraffine, the immersion in the heated paraffine being made as short as possible, usually not more than fifteen minutes. Sections about  $5\mu$  thick were then cut on a Minot microtome.

For staining, Heidenhain's iron haematoxylin was employed almost exclusively. Other stains were tried, but none gave such satisfactory results.

## II. DESCRIPTION OF TESTIS.

The paired testes lie on the ventral side of the abdominal cavity, one on each side of the alimentary canal. Each adult testis is a flattened pear-shaped body, consisting of seven approximately equal lobes arranged in a single row and with no indication of a division into groups, as in *Pyrrhochoris* (Mayer, '74), or of a difference in size, as in *Pentatoma* (Montgomery, '98). During the first few instars the testis is white, but after the third moult a faint yellow color makes its appearance toward the free end. This gradually deepens to orange, and then just before the fifth or last moult, or in some cases earlier, — different individuals varying in this respect, — a red pigment appears between the lobes and spreads out in streaks across them. By the time the insect emerges from its hibernation the red pig-