

**BIOLOGICAL LECTURES
DELIVERED AT THE MARINE
BIOLOGICAL LABORATORY
OF WOODS HOLL 1899**

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649074365

Biological Lectures Delivered at the Marine Biological Laboratory of Woods Holl 1899 by
Marine Biological Laboratory

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd.
Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

MARINE BIOLOGICAL LABORATORY

**BIOLOGICAL LECTURES
DELIVERED AT THE MARINE
BIOLOGICAL LABORATORY
OF WOODS HOLL 1899**

BIOLOGICAL LECTURES

FROM LIBRARY

THE MARINE BIOLOGICAL LABORATORY
OF WOODS HOLL

1899



BOSTON, U.S.A.
GINN & COMPANY, PUBLISHERS
The Athenæum Press
1900

4

YVA98L1 3MA1

COPYRIGHT, 1900
By GINN & COMPANY
ALL RIGHTS RESERVED

2511
M33
1899

CONTENTS.

LECTURE		PAGE
I.	<i>The Evolution of the Sporophyte in the Higher Plants.</i> DOUGLAS HOUGHTON CAMPBELL . . .	1
II.	<i>The Nature of the Evidence exhibited by Fossil Plants, and its Bearing upon our Knowledge of the History of Plant Life.</i> D. P. PENHALLOW . . .	19
III.	<i>Influence of Inversions of Temperature, Ascending and Descending Currents of Air, upon Distribution.</i> PROF. D. T. MACDOUGAL . . .	37
IV.	<i>Significance of Mycorrhizas.</i> PROF. D. T. MACDOUGAL	49
V.	<i>Instinct.</i> EDWARD THORNDIKE	57
VI.	<i>The Associative Processes in Animals.</i> EDWARD THORNDIKE	69
VII.	<i>The Behavior of Unicellular Organisms.</i> HERBERT S. JENNINGS	93
VIII.	<i>The Blind-Fishes.</i> CARL H. EIGENMANN	113
IX.	<i>Some Governing Factors usually neglected in Biological Investigations.</i> ALPHEUS HYATT	127
X.	<i>On the Development of Color in Moths and Butterflies.</i> ALFRED GOLDSBOROUGH MAYER	157
XI.	<i>The Physiology of Secretion.</i> A. MATHEWS	165
XII.	<i>Regeneration: Old and New Interpretations.</i> T. H. MORGAN	185
XIII.	<i>Nuclear Division in Protozoa.</i> GARY N. CALKINS	209
XIV.	<i>The Significance of the Spiral Type of Cleavage and its Relation to the Process of Differentiation.</i> C. M. CHILD	231
XV.	<i>The Aims of the Quantitative Study of Variation.</i> C. B. DAVENPORT	267
XVI.	<i>On the Nature of the Process of Fertilisation.</i> JACQUES LOEB	273

FIRST LECTURE.

THE EVOLUTION OF THE SPOROPHYTE IN THE HIGHER PLANTS.

DOUGLAS HOUGHTON CAMPBELL.

THE questions relating to the origin of organic structures must always possess great interest for the biologist, and when I was asked to speak before the students at Woods Holl, it seemed to me that a discussion of recent views bearing on the development of the spore-producing structures of the higher plants, *i.e.*, the Archegoniates and seed-bearing plants, would not be inappropriate.

There is good reason to suppose that among plants, as among animals, the most primitive forms were aquatic; and it is highly probable that many existing fresh-water algæ are but slightly modified descendants of these ancestral types. This is evinced by the great uniformity shown by existing green algæ all over the world. Most genera and many species are cosmopolitan, and they exhibit far less variety than is shown by their larger and more specialized marine relations. Presumably the conditions in fresh water have changed but little, and as the more specialized forms have taken to the land, or have developed in the sea, the few that remained in their primitive environment have been subjected to much less competition, and have probably persisted with but little change from very remote times.

Leaving aside certain forms of doubtful affinities, like the bacteria and blue-green algæ, the existing forms which represent most nearly the ancestors of the higher plants are the Volvocineæ and Protococcaceæ. The former are, as all botanists know, free-swimming green organisms which show resem-

blances on the one hand to low animals, and on the other are very like the free-swimming reproductive cells, or zoöspores, of many of the higher algæ. The latter probably originated from forms like the simpler Volvocineæ by the loss of motility associated with the development of a continuous cellulose membrane about the vegetative cells. This stage in the evolution of the green algæ is represented by some of the simpler Protococcaceæ. Above these unicellular forms are a number of filamentous plants, single rows of nearly uniform cells. Such plants are *Œdogonium* or *Conferva*, which well represent the next step in the evolution of the plant body.

The differentiation of the reproductive cells in the algæ, while it offers one of the most interesting and instructive examples of the evolution of plant structures, must be passed over here. It may be mentioned, however, that the differentiation of the sexual cells has evidently taken place quite independently in several groups of algæ.

In the further development of plant types two very important factors are to be considered. First, the adaptation to a marine existence, and, second, the exchange of an aquatic for a terrestrial life.

The conditions in the ocean are markedly different from those in fresh water, and most plants which have adapted themselves to life in the sea have become decidedly changed. The salinity of the water has no doubt been one of the factors in these changes, but more important is probably the question of light. The two most characteristic groups of sea plants, the red and brown seaweeds, are provided with special pigments in addition to the chlorophyll, and there is little question that these pigments are developed, in part at least, in response to changed light conditions.

As the conditions of light and temperature in a marine environment are far more constant than those in fresh water, we find, as a rule, that marine algæ, especially those inhabiting the deeper water, are more susceptible to changes of light and temperature than are most fresh-water forms.

While certain seaweeds growing between tide-marks are subject to exposure to the air, it is only for a short time, and

we find in most such algæ a development of mucilaginous tissue which prevents their complete desiccation.

These seaweeds have adapted themselves perfectly to their peculiar environment, and such highly specialized forms as the great kelps and many red algæ probably represent the highest types of these marine plants. They have diverged widely from the simpler fresh-water algæ, and there is no reason to suppose that they have given rise to any higher types.

The simple fresh-water green algæ, which, so far as we know, most nearly represent the ancestors of the terrestrial green plants, differ much in their conditions of life from the seaweeds. Most bodies of fresh water are subject to great fluctuations of depth, often drying up completely for long periods, or sometimes being frozen. It is obvious that plants living under such conditions must be very resistant, and we find that such is the case among most green algæ. Not only, as a rule, are they capable of enduring a great range of temperature, but usually at the end of their vegetative period they produce special cells, "spores," which can endure complete desiccation without injury, and are also uninjured by freezing. By means of these "resting-spores" the plant is carried over from one growing period to the next, and when the conditions are favorable, the spores germinate and give rise to a new generation. This production of resting-spores is one of the most striking differences between these fresh-water algæ and their red and brown relations in the sea, where there is usually no necessity for such resting-spores.

Certain green algæ, like some species of the common genus *Vaucheria*, may be considered amphibious, as they do not actually grow in the water, but are exposed to the air on the surface of moist earth, from which they absorb the water necessary for their growth. The ability to thus grow with a diminished water supply is an evident advantage, and in some such way as this it is probable that the first strictly terrestrial plants originated from some originally aquatic algal ancestors.

We can imagine some such forms gradually becoming better and better able to vegetate on the mud left by the subsidence of the water, and finally becoming adapted to life on land. The