

THE MEDICAL TREATMENT OF GALL-STONES

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649645503

The Medical Treatment of Gall-Stones by J. H. Keay

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd.
Cover @ 2017


This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

J. H. KEAY

**THE MEDICAL
TREATMENT OF
GALL-STONES**

THE MEDICAL TREATMENT
OF
GALL-STONES



THE MEDICAL TREATMENT
OF
GALL-STONES

BY
J. H. KEAY, M.A., M.D.



PHILADELPHIA
P. BLAKISTON'S SON & CO.
1012 WALNUT STREET
1902

Printed in England

HARVARD UNIVERSITY
SCHOOL OF MEDICINE AND PUBLIC HEALTH

LIBRARY
Gift: Mrs. R. W. Wilcox
27 FEB 1937

A 16.K.1902.1

Wh:

Desc
c
(
a
tl
(
n
(
p
(
di

CONTENTS

INTRODUCTORY

	PAGES
Why write about gall-stones?—Reasons given—Before resorting to operation consider (1) that the mere presence of gall-stones affords no reason for operation; (2) that, of those who suffer, more are cured by medical and hygienic means than surgical; (3) that operation may relieve symptoms, but does not radically cure the disease	1—10

CHAPTER I

THE FORMATION OF GALL-STONES

Description of gall-stones—Cause of formation readily discovered—Materials of which gall-stones are composed: (1) cholesterin, (2) lime (bilirubinate of lime)—Cholesterin and lime chiefly derived from gall-bladder and bile-ducts, their excretion in more than normal quantity being due to (1) the irritating properties of bile developed during stagnation, or (2) derived from the products of indigestion, (3) bacteria—The formation of gall-stones indirectly dependent on (1) heredity, (2) age, (3) sex, (4) occupation, (5) locality, (6) diet and digestion, (7) worry, (8) other diseases	11—34
---	-------

CHAPTER II

THE MIGRATION OF GALL-STONES

PAGES

Gall-stones give no trouble when at rest in gall-bladder—Their migration seldom caused by (1) a fall or jerk, but almost always by (2) the development of bacteria or additional irritants in the bile—When set in motion they (a) pass directly through the ducts into the duodenum; or (b) become impacted in the ducts—jaundice; or (c) perforate bladder or ducts—fistulæ	35—48
---	-------

CHAPTER III

MORBID CONDITIONS ASSOCIATED WITH GALL-STONES

Morbid conditions described—Pathological changes in gall-bladder or bile-ducts not always associated with, and seldom due to, the mechanical irritation of stones—The formation of stones and alteration in form or structure of gall-bladder or bile-ducts due to certain inflammatory conditions—Pathological changes in gall-bladder: contraction, simple catarrh, dropsy, gangrene, empyæma, adhesions, cancer	49—69
--	-------

CHAPTER IV

SYMPTOMS AND DIAGNOSIS

Importance of early diagnosis—Symptoms and signs: (1) Pain, caused by (a) the passage of stones through the ducts, (b) an inflamed or distended gall-bladder, (c) adhesions; (2) tenderness; (3) the evacuation of biliary calculi; (4) jaundice; (5) fever; (6) vomiting; (7) disordered appetite; (8) general bodily condition—Differential diagnosis: (1) Cramp of stomach; (2) gastric ulcer; (3) appendicitis; (4) wandering kidneys; (5) cystic kidney, ovarian cysts, etc.; (6) pancreatic cysts, calculi, etc.; (7) cancer; (8) lead colic—Methods of diagnosis	70—93
---	-------

CONTENTS

vii

CHAPTER V

TREATMENT

PAGES

Surgical <i>v.</i> medical and hygienic treatment ; medical treatment the more radical—The chief aim of medical treatment is not (a) to expel stones, nor (b) to dissolve them, but to prevent their formation and to keep them at rest by (1) removing those conditions that give rise to deterioration of bile and consequent irritation of gall-bladder and bile-ducts— <i>e.g.</i> , errors of digestion, worry, constriction by dress, etc.—and by (2) the use of drugs, mineral waters, etc., which promote a free flow of healthy bile—Narcotics and sedatives—Drugs that promote flow of healthy bile—Turpentine and ether—Olive-oil—Mineral waters—Surgical treatment—Operation objected to in certain conditions—Who is to operate?—Indications for operation given by various surgeons—Those given by Mayo Robson discussed—Conclusions	94—126
--	--------